



TECHNICAL INFORMATION BULLETIN
FOR THE USE OF Venerate XC® BIOINSECTICIDE
First Generation Codling Moth Larvae on Pome Fruit

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR USE IN ALL STATES WHERE CROP AND PEST(S) EXIST AND WHERE 2(Ee) RECOMMENDATIONS ARE RECOGNIZED.

Venerate XC®
EPA Reg. No. 84059-14
DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Crop	Target Insect	Application Method	Product Use Rate per Application	Application Instructions
Pome fruit (apples, crabapple, loquat, mayhaw, pears and quince)	First generation codling moth larvae (<i>Cydia pomonella</i>)	Foliar (Ground application) in a minimum of 50 gallons of water per acre	1-2 quarts per acre	<p>Careful attention to the codling moth cycle is required for successful control. Closely follow regional spray recommendations for codling moth based upon biofixes, pheromone trap catches and egg hatch.</p> <p>Codling moth overwinter as mature larvae in silken cocoons, with adults emerging around bloom and proceeding to lay eggs. There are two to three generations per season. Applications should be timed for egg hatch of first generation larvae for maximum control. Codling moth larvae must be controlled before they penetrate developing fruit.</p> <p>Venerate can be used in combination with mating disruption programs.</p>
<p>Venerate XC® has a pre-harvest interval (PHI) of 0 days. Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.</p>				

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

This recommendation must be in the possession of the user at the time of pesticide application.

Venerate XC® is a registered trademark of Marrone Bio Innovations, Inc.
© Copyright Marrone Bio Innovations, Inc., 2020
Marrone Bio Innovations, Inc.
1540 Drew Ave.
Davis, California 95618
Telephone (530) 750-2800 www.marronebioinnovations.com