



TECHNICAL INFORMATION BULLETIN FOR THE USE OF GRANDEVO® TO CONTROL CODLING MOTH ON TREE NUTS INCLUDING WALNUTS

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR USE IN ALL STATES WHERE CROP AND PEST(S) EXIST AND WHERE 2(EE) RECOMMENDATIONS ARE RECOGNIZED.

Grandevo[®] EPA Reg. No. 84059-17

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CROP DISEASE RATE	USE RECOMMENDATIONS
Tree nuts including almonds, cashew, chestnut, filbert (hazelnut), macadamia nut, pistachios and walnut Tree nuts (Cydia (Cydia pomonella)) Codling moth (Cydia Grandevo per acre when used alone, or 1-3 pounds per acre when tank mixed with another insecticide labeled for control of codling moth. Use the higher rate range for larger trees and heavier infestations. Grandevo per acre when codling moth used alone, or 1-3 pounds codling per acre when tank mixed with another insecticide labeled for control of codling moth. Grandevo per acre when used alone, or 1-3 pounds codling per acre when tank mixed with another insecticide labeled for control of codling moth. Grandevo per acre when used alone, or 1-3 pounds codling per acre when tank mixed with another insecticide labeled for control of codling moth. Grandevo per acre when used alone, or 1-3 pounds codling per acre when tank mixed with another insecticide labeled for control of codling moth. Grandevo per acre when used alone, or 1-3 pounds per acre when tank mixed with another insecticide labeled for control of codling moth. Grandevo per acre when used alone, or 1-3 pounds per acre when tank mixed with another insecticide labeled for control of codling moth. Grandevo per acre when tank mixed with another insecticide labeled for control of codling moth. Grandevo per acre when tank mixed with another insecticide labeled for control of codling moth.	Apply Grandevo as a foliar spray when pests first appear. Closely follow regional spray recommendations for codling moth based upon biofixes, pheromone trap catches and egg hatch. Grandevo is a larvicide and must be applied prior to egg natch. Codling moth larvae must be controlled before hey penetrate developing nuts. Codling moth egg hatch ypically begins at 220 – 250 degree days (base 50° F). Reapply on a 3-7 day interval for the duration of egg natch. Grandevo can be used in combination with mating disruption programs.

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

This recommendation must be in the possession of the user at the time of pesticide application.

Grandevo[®] is a registered trademark of Marrone Bio Innovations, Inc. © Copyright Marrone Bio Innovations, Inc., 2013

Marrone Bio Innovations, Inc. 2121 Second Street, Suite B-107 Davis, California 95618 Telephone (530) 750-2800 www.marronebioinnovations.com

Phone: 530-750-2800